



JUL 2

<b>REPORT DOCUMENTATION PAGE</b>			Form Approved OMB No. 0704-0188	
<b>AD-A267 085</b>			<small>It is requested to average 1 hour per reference, including the time for reviewing instructions, searching existing data sources, gathering and reviewing the collection of information, time for reviewing existing data sources, and reviewing the collection of information. Some comments regarding the burden estimate of this and the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, Suite 1204, Washington, DC 20540-6188.</small>	
			<b>REPORT DATE</b> June 23, 1993	
			<b>REPORT TYPE AND DATES COVERED</b> Reprint	
<b>4. TITLE AND SUBTITLE</b> Dynamic potential barrier effects in hydrogen tunneling in trans-cis isomerizations			<b>5. FUNDING NUMBERS</b> 61102F 2303/B3	
<b>6. AUTHOR(S)</b> Zhongzhi Yang, Yue Qin, and Donald L. Thompson				
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> Oklahoma State University Stillwater, OK 74078  AFOSR-TR			<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>  93 05 29	
<b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> AFOSR/NC 110 Duncan Ave., Suite B115 Bolling AFB, DC 20332-0001			<b>10. SPONSORING/MONITORING AGENCY REPORT NUMBER</b>  AFOSR-90-0018	
<b>11. SUPPLEMENTARY NOTES</b> Chemical Physics Letters, Vol. 206, No. 5,6, 7 May 93 pp. 445-449				
<b>12a. DISTRIBUTION/AVAILABILITY STATEMENT</b> APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED.			<b>12b. DISTRIBUTION CODE</b> 	
<b>13. ABSTRACT (Maximum 200 words)</b>  <p>A simple model is employed to illustrate the influence of the dynamic effective potential resulting from torsional motion on tunneling via bending modes in trans-cis isomerization in polyatomic molecules. We show that the effect is significant and can determine the dominant tunneling mode in systems such as HSiOH.</p>				
<b>14. SUBJECT TERMS</b> Chemical dynamics, Tunneling Effects,			<b>15. NUMBER OF PAGES</b> 5	
			<b>16. PRICE CODE</b>	
<b>17. SECURITY CLASSIFICATION OF REPORT</b> UNCLASSIFIED	<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> UNCLASSIFIED	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> UNCLASSIFIED	<b>20. LIMITATION OF ABSTRACT</b>	

## Dynamic potential barrier effects in hydrogen tunneling in trans-cis isomerizations

Zhongzhi Yang<sup>1</sup>, Yue Qin<sup>2</sup> and Donald L. Thompson

*Department of Chemistry, Oklahoma State University, Stillwater, OK 74078, U.S.A.*

Received 17 December 1992; in final form 15 February 1993

A simple model is employed to illustrate the influence of the dynamic effective potential resulting from torsional motion on tunneling via bending modes in trans-cis isomerization in polyatomic molecules. We show that the effect is significant and can determine the dominant tunneling mode in systems such as HSiOH.

### 1. Introduction

Most treatments of barrier tunneling make use of simplifying assumptions, such as one-dimensional paths. As chemists, we are often interested in processes that occur in polyatomic molecules, thus the tunneling processes of interest are often multi-dimensional. Makri and Miller [1] have presented a method for treating tunneling in multi-dimensional systems semiclassically. However, it is often necessary and appropriate to calculate tunneling corrections by assuming that it occurs along some special "tunneling direction", for example, along a normal mode direction.

An example where this approach may be reasonable is trans-cis isomerization. It is often a good approximation to treat the torsional motion as separable. However, as *ab initio* calculations have shown, trans-cis isomerization can occur via an angle bending mode in some molecules. The purpose of this Letter is to point out that there are dynamic effects which arise from the torsional motion that must be taken into account for tunneling along a bending mode direction. We use a very simple model to illustrate the qualitative effects of the torsional mo-

tion on the bending mode tunneling. We consider the simplest case, that is, a four-atom molecule, namely, HSiOH.

The cis and trans conformers of HSiOH, and its isotopic (D and <sup>18</sup>O) analogs, have received a lot of attention lately. Ismail et al. [2] studied the interaction of silicon atoms and water molecules in a solid argon matrix. Infrared spectra of the products (including cis and trans conformers of HSiOH, HSi<sup>18</sup>OH, HSiOD, and HSi<sup>18</sup>OD) were measured and assigned. Withnall and Andrews [3] studied the matrix reactions of silane and oxygen atoms and identified HSiOH, HSi<sup>18</sup>OH, DSiOD, and DSi<sup>18</sup>OD by infrared absorption. Kudo and Nagase [4,5] have reported *ab initio* calculations for H<sub>2</sub>SiO and related molecules, including trans- and cis-HSiOH. They found that there are two possibilities for the trans to cis isomerization transition state: either along an in-plane bending mode or along the torsional direction. Their results showed that the latter, which has a barrier of 11 kcal/mol, is 1.1 kcal/mol lower than the former. Sakai and Jordan [6] have reported SCF results for the structures, stabilities, and normal mode frequencies for cis- and trans-HSiOH. (They also reported results for similar molecules. They found that the lowest energy path for trans→cis isomerization in HAlOH involves the in-plane motion of the H-atom bonded to the O-atom rather than the torsional motion as in the case of HBOH, HCOH, and HSiOH.) Tachibana et al. [7,8] have investigated

<sup>1</sup> Permanent address: Institute of Theoretical Chemistry, Jilin University, Changchun, 130023, People's Republic of China.

<sup>2</sup> Current address: Department of Chemistry, University of Oklahoma, Norman, OK, USA.

tunneling in the cis-trans isomerization in HSiOH by using an ab initio IRC (intrinsic reaction coordinate) method. They proposed that the isomerization path may be regarded as one-dimensional along the direction of the SiOH bending mode. That is, the path maintains  $C_s$  symmetry and the vibrational modes orthogonal to it do not contribute to the dynamics along the path.

We should also point out some classical dynamics studies that are relevant to this problem. Spears and Hutchinson [9,10] used classical mechanics to study the dynamics of vibrationally excited trans-diimide. Using a very simple model and classical trajectory calculations, they showed that the isomerization dynamics of diimide are dominated by the "fictitious" forces arising from the presence of torsional motion and the source of the final "kick" of energy into the isomerization coordinate (torsion) arises from a Coriolis coupling of the torsion to the NH stretches. They pointed out that the isomerization via a planar bending mechanism is prevented since such motion entails crossing an infinite effective potential due to the centrifugal force. That is, they concluded that the bending pathway is inactive.

We focus here on the dynamic effects on tunneling paths between the trans and cis conformers of molecules of the type HABC, taking HSiOH as an example. We use a simple model to illustrate the effect of the dynamic effective potential, which arises from the torsional motion, on the tunneling probability. It is pointed out that this "dynamic" correction to the "static" potential in one-dimensional treatments of tunneling for cis-trans isomerizations may be important. We also note that due to the existence of this dynamic effect, the realistic tunneling path may be a curved route that is a combination of the torsional and bending modes, and is possibly coupled with the stretching modes as well.

## 2. The dynamic effective potential barrier

The Born-Oppenheimer approximation allows the Schrödinger equation for a molecule to be separated into two equations,

$$(T_e + V_{ee} + V_{en} + V_{nn})\Psi_e(\mathbf{R}, \mathbf{r}) = E(\mathbf{R})\Psi_e(\mathbf{R}, \mathbf{r}), \quad (1)$$

$$[T_n + E(\mathbf{R})]\Psi_n(\mathbf{R}) = \epsilon\Psi_n(\mathbf{R}), \quad (2)$$

where  $\Psi_e(\mathbf{R}, \mathbf{r})$  and  $\Psi_n(\mathbf{R})$  are the electronic and nuclear wavefunctions, respectively.  $\mathbf{R}$  and  $\mathbf{r}$  are the position vectors of the nuclei and electrons, respectively;  $T_e$ ,  $T_n$ ,  $V_{ee}$ ,  $V_{en}$ , and  $V_{nn}$  are the operators for the kinetic energies for the nuclei and electrons, electron repulsions, electron-nuclear attractions, and nuclear repulsions, respectively;  $\epsilon$  is the total energy and  $E(\mathbf{R})$  is the "potential energy surface" (which we refer to as the static potential).

Now, let us consider the trans-cis isomerization in HABC-type molecules taking HSiOH as an example. The ab initio calculations [4,7] show that the trans- and cis-HSiOH conformers exist in local minima with planar structures on the ground-state potential energy surface. The two minima are separated by barriers of about 10 kcal/mol along both the out-of-plane torsional coordinate and the in-plane bending mode direction. Kudo and Nagase [4] and Sakai and Jordan [6] have suggested that the torsional mode is the dominant pathway for the isomerization. However, Tachibana et al. [7,8] assumed that the isomerization involves pure SiOH bending motion and computed tunneling probabilities for passage through the barrier along that coordinate. The dominant pathway for the isomerization is determined simply by the relative heights of the two barriers on the static potential energy surface. In the following, we just show that the effective potential, which arises from dynamic effects, should be considered for this process.

The coordinate system for our model is illustrated in fig. 1. The SiO bond is along the  $z$  axis with the O-atom at the origin. Points A, B, C, D, and E are all located on a sphere of radius  $R_0$ , the equilibrium OH bond length. The equilibrium cis and trans geometries both lie in the YOZ plane. The OH bond points to the upper right for the trans geometry and the upper left for the cis. As the molecular translation and rotation are separated out, the static potential can be written generally as  $V(r_1, r_2, \theta_1, \theta_2, \varphi_1 - \varphi_2)$ , where  $r_1$ ,  $\theta_1$ , and  $\varphi_1$  describe the first H-atom, and  $r_2$ ,  $\theta_2$ , and  $\varphi_2$  the second. The torsional potential depends only on the difference  $\varphi_1 - \varphi_2$ .



along the reaction coordinate  $s$ ,  $V_1(s, x) = 0$  and  $K(s)x = 0$ , which leads to eq. (7).

It should be noted that the case we treat, the effect of torsional motion on the tunneling path, is different to those usually considered [11,12]. The usual treatment is for vibrational modes that are normal to the path  $s$  with a harmonic or modified-harmonic model, and where the vibrational energy only about the path  $s$  is considered.

Let us now make a rough estimate of the effect of the dynamic effective potential on the tunneling path in the trans  $\rightarrow$  cis isomerization of HSiOH. We refer again to fig. 1, which illustrates isomerization along the pathways with torsional motion. When the hydrogen atom is at point A  $\tau = 0$  and  $\theta = 65.5^\circ$ , at point B  $\tau$  is undefined and  $\theta = 0^\circ$ , at point C  $\tau = 180^\circ$  and  $\theta = 65.5^\circ$ , at point D  $\tau$  is between  $0^\circ$  and  $90^\circ$  and  $\theta = 65.5^\circ$ , and at point E  $\tau = 90^\circ$  and  $\theta < 65.5^\circ$ . Pure planar isomerization is along the path ABC. The more general path involves a torsion plus bend motion as illustrated by the points ADEC in fig. 1, where it is visualized as being broken down into simple segments of the two motions. We can roughly estimate this path by assuming it follows the smallest polar angle  $\theta = (65.5^\circ/90^\circ) \tau$  as the atom moves from A to C. Quantum mechanically, the internal rotation operator  $L^2$  has an average deviation (uncertainty) on the order of  $\hbar^2$ . Thus, the dependence of the largest value of the dynamic effective potential along the path on  $\tau$  can be expressed approximately as

$$V_{\text{dep}}(\tau) \sim \hbar^2 \left[ 2\mu R_0^2 \sin^2\left(\frac{65.5^\circ}{90^\circ} \tau\right) \right]^{-1} \\ = 0.05 \sin^{-2}(0.73\tau) \text{ kcal/mol.} \quad (8)$$

The cis and trans conformers are at about the same energy and the barriers for the two isomerization channels on the static potential energy surface are roughly at the same level (about 10 kcal/mol above the conformer wells). We approximate the barrier height of the total effective potential as

$$V_0(\tau) = [10.0 + 0.05 \sin^{-2}(0.73\tau)] \text{ kcal/mol,} \quad (9)$$

in order to simply show the dynamic effect. The dependence of  $V_0(\tau)$  on  $\tau$  is plotted in fig. 2. It decreases rapidly as the torsional angle increases.




Fig. 2. The barrier height of the approximate effective potential as a function of the torsional angle  $\tau$ , eq. (9).

### 3. Tunneling probability

To estimate the effect of the dynamic effective potential on the tunneling, we represent the potential with the Eckart barrier [13].

$$V(s) = V_0(\tau) \text{sech}^2(s), \quad (10)$$

where  $s$  is the tunneling coordinate. Fig. 3 shows the shapes of the potential barrier for a few  $\tau$ , where the lowest one is the static potential.

The one-dimensional Schrödinger equation along the tunneling path,

$$\frac{d^2\psi(s)}{ds^2} + \frac{2m}{\hbar^2} [E - V_0 \text{sech}^2(s)] \psi(s) = 0, \quad (11)$$




Fig. 3. The Eckart barriers used to represent the static potential and the dynamic effective potentials for torsional angles  $\tau = 5^\circ$ ,  $10^\circ$ ,  $15^\circ$ , and  $20^\circ$ .

can be solved analytically [13]. The permeability is

$$P_{\text{tun}} = \sinh^2(\pi k) \times [\sinh^2(\pi k) + \cosh^2 \sqrt{\frac{1}{2} \pi (8mV_0/\hbar^2 - 1)}]^{-1}, \quad (12)$$

where

$$k = \sqrt{2mE/\hbar}.$$

The dependence of tunneling probability  $P_{\text{tun}}$  on the torsional angle  $\tau$  for  $E=8$  kcal/mol is shown in fig. 4. The probability of tunneling is zero at  $\tau=0$  since the total effective potential is infinite. As  $\tau$  increases  $V_{\text{dep}}$  decreases rapidly (see fig. 2) and the tunneling probability becomes significant at about  $\tau=20^\circ$ .

The torsional energy levels are, of course, quantized with relatively large spacings. We have not considered the quantized levels here, although one could easily do so. We merely wish to illustrate the effect and thus have considered the trans-cis motion on a potential energy surface where the torsional mode might be considered to be any state, generally, not an eigenstate of the torsion.

We have ignored the effects of the variation in the width of the barrier, which is certainly significant. We have assumed that all paths have the same width since we are interested in merely illustrating the qualitative dynamic effect of the torsional motion on the tunneling paths. Work is in progress on a more accurate and elaborate treatment of this problem.




Fig. 4 The tunneling probability as a function of the torsional angle.

#### 4. Conclusions

We have used a simple model to illustrate the effect of the dynamic effective potential that results from torsional motion on tunneling paths along the trans-cis isomerization coordinate. This effect may be important in many systems and must be explicitly considered in one-dimensional treatments of tunneling. Because of this effect, the torsional or a curved torsional-plus-bend pathway, possibly coupled to stretching, is likely to be preferred over the pure in-plane bending pathway.

#### Acknowledgement

ZY appreciates the kindness and assistance of Dr. Karen Bintz, Ms. Candee Chambers, and Mr. Paul Zahner during his visit. This work was supported by the Air Force Office of Scientific Research under Grant AFOSR-90-0048.

#### References

- [1] N. Makri and W.H. Miller, *J. Chem. Phys.* 91 (1989) 4026.
- [2] Z.K. Ismail, R.H. Hauge, L. Fredin, J.W. Kauffman and J.L. Margrave, *J. Chem. Phys.* 77 (1982) 1617.
- [3] R. Withnall and L. Andrews, *J. Phys. Chem.* 89 (1985) 3261.
- [4] T. Kudo and S. Nagase, *J. Phys. Chem.* 88 (1984) 2833.
- [5] T. Kudo and S. Nagase, *Chem. Phys. Letters* 128 (1986) 507.
- [6] S. Sakai and K.D. Jordan, *Chem. Phys. Letters* 130 (1986) 103.
- [7] A. Tachibana, H. Fueno, M. Koizumi, T. Yamabe and K. Fukui, *J. Phys. Chem.* 92 (1988) 935.
- [8] A. Tachibana, H. Fueno and T. Yamabe, *J. Am. Chem. Soc.* 108 (1986) 4346.
- [9] L.G. Spears Jr. and J.S. Hutchinson, *J. Chem. Phys.* 88 (1988) 240.
- [10] L.G. Spears Jr. and J.S. Hutchinson, *J. Chem. Phys.* 88 (1988) 250.
- [11] R.A. Marcus, *J. Chem. Phys.* 45 (1986) 4493.
- [12] W.H. Miller, N.C. Handy and J.E. Adams, *J. Chem. Phys.* 72 (1980) 99.
- [13] W.H. Miller, *J. Phys. Chem.* 87 (1983) 3811.
- [13] R.P. Bell, *The tunnel effect in chemistry* (Chapman and Hall, London, 1980).

DTIC QUALITY INSPECTED 1

449

Special  
A-1/20